THE DISK SETTLING INSTABILITY


Fast clumping of small grains in protoplanetary disks

J. Squire — P. Hopkins arXiv:1711.03975



ABSTRACT

We identify and characterize the disk "settling instability," a variant of the streaming instability (Youdin & Goodman 2005) that can occur in protoplanetary disks when grains settle vertically into the midplane region. The growth rate of the settling instability is independent of grain size and comparable to the disk orbital frequency for realistic parameters, suggesting that the instability is relevant for even the smallest grains. In addition, its characteristic wavelengths are larger than those of the streaming instability, potentially allowing the instability to concentrate large masses of solids. These properties suggest that in the process of settling, small grains could band into rings (then filaments or clumps) with significantly higher than background metallicity. This could have a variety of important consequences for the early stages of planetesimal formation, for instance, by enhancing coagulation rates of small grains and/or creating high-metallicity regions in the midplane that act to seed the standard streaming instability.




RESONANT DRAG INSTABILITY arXiv:1706.05020

An algorithm to find instabilities in coupled dust dust-gas mixtures

- 1) Work out equilibrium: drift velocity (ws), gas properties
- 2) Choose a gas wave, with frequency $\omega_{\text{wave}}(\boldsymbol{k})$



- 3) Match the dust frequency $k \cdot \mathbf{w}_s$ to $\omega_{\text{wave}}(k)$ gives "resonant wavenumber" $k_{\rm res}$
- 4) Dust "sees" a stationary wave, clumps dust dust backreaction on gas drag forces on dust Growth rate easy to calculate $\approx \mu^{1/2} \left[(\xi_{\mathcal{F}}^L T_{\rho_d}^{(1)}) (\mathbf{k}^T D_{\mathrm{drag}}^{-1} C_{\mathbf{v}} \xi_{\mathcal{F}}^R) \right]^{1/2}$



 \leftarrow long wavelengths $k\eta r$

COUPLED GAS+DUST EQUATIONS IN A LOCAL PATCH OF DISK

Dust in midplane, streams through gas with velocity

$$\mathbf{w}_s = \mathbf{v}_{\mathrm{dust}} - \mathbf{u}_{\mathrm{gas}} = -2 \frac{\eta \, U_K (1 + \mu) \, \tau_s}{(1 + \mu)^2 + \tau_s^2} \, \hat{\mathbf{x}} + \frac{\eta \, U_K \tau_s^2}{(1 + \mu)^2 + \tau_s^2} \, \hat{\mathbf{y}}$$
 $\approx -2 \eta U_K \tau_s \, \hat{\mathbf{x}}, \quad \mu \ll 1, \, \tau_s \ll 1 \, \text{(Nakagawa-Sekiya-Hayashi drift)}$

Gas epicycles have frequency $\omega_{\rm gas} = \pm \frac{\kappa_z}{k} \Omega$


occurs at

Resonance (fastest growing mode)


$$k_{\rm res} = \frac{\Omega}{|\mathbf{w}_{s,r} \tan \theta_{k}|}$$

Growth rate is $\Im(\omega) \approx \Omega \, \tau_s \sqrt{\frac{\mu}{8}} (2\hat{k}_z^2 + \hat{k}_x^2)$

STREAMING INSTABILITY




Occurs because the streaming dust excites gas epicycles



Growth rate $\theta_k = 30^{\circ}$

At $\mu > 1$ a different mode dominates (see appendix of paper) Growth rate $\omega \approx i\sqrt{\mu-1}$, slower (but larger scales) for large grains

Dust also settles vertically if spread throughout disk



Resonates with gas epicycles $\omega_{\rm gas} = \pm \frac{k_z}{l_z} \Omega$ than radial drift


Resonance (fastest growing mode) occurs at


$$k_{\rm res} = \frac{\Omega}{|\tan \theta_k \mathbf{w}_{s,x} + \mathbf{w}_{s,z}|} \approx \frac{\Omega}{|\mathbf{w}_{s,z}|} \sim 30 \times \text{larger scale than}$$
 streaming instability

Growth rate is

$$\Im(\omega) \approx \Omega\left(\frac{\mu}{2}\right)^{1/2} |\sin \theta_k|$$
 Growth rate does not depend on grain size!


SETTLING INSTABILITY





Weak dependence on metallicity ($\sim \mu^{1/2}$)


 \leftarrow long wavelengths $\log_{10}(k_x\eta r)$



"Double-resonant mode"

For mode angle $\tan \theta_k = w_{s,z}/w_{s,x}$

Minor modifications from gas stratification see also Lambrechts+ (2016)



Grows faster than settling for $\tau_s \lesssim 0.1 \left(\frac{\mu}{0.01}\right)^{1/2}$

Grains sizes not affected by gas viscosity $R_d \gtrsim 8 \left(\frac{r}{\text{AU}}\right)^{-3/4} \mu\text{m}$

May be affected by turbulence for smaller grains $\tau_s \lesssim 0.01 \left(\frac{\alpha}{10^{-4}}\right)^{1/2} \left(\frac{\mu}{0.01}\right)^{-3/4}$

(but generally less so than YG streaming instability)

Enough mass in radial wavelength for planetesimal of size $R \sim 1000 \,\mathrm{km} \, \left(\frac{\tau_s}{0.001}\right)^{1/3} \, \left(\frac{\mu}{0.01}\right)^{1/3} \, \left(\frac{r}{\mathrm{AII}}\right)^{-0.26}$

Possible IMPLICATIONS


Causes local density

enhancements

Clumps contain a large

mass of solids

For smaller grains, growth time much shorter than settling time $t_{\rm settle} \sim (\Omega \tau_s)^{-1}$



DUST

Enhances coagulation rate in clumps during settling? Drążkowska & Dullemond (2014)

Would naturally produce highmetallicity regions in midplane

Simulations show a critical metallicity $Z \gtrsim 3Z_{MMSN}$ for the streaming instability to form planetesimals (Johansen+ 2009, Bai & Stone 2010, Carrera+ 2015)

Does the disk settling instability provide high-Z seed regions for the YG streaming instability?

Or enhanced densities could seed pressure bumps?

Or even direct collapse of high-density regions?

Numerical simulations needed...